

**BRITISH ZOOPHYTES.**

**VOL. II.**

A  
HISTORY  
OF THE  
BRITISH ZOOPHYTES.

BY  
GEORGE JOHNSTON, M.D., LL.D.,  
FELLOW OF THE ROYAL COLLEGE OF SURGEONS OF EDINBURGH.



IN TWO VOLUMES.—VOL. II.

SECOND EDITION.

LONDON:  
JOHN VAN VOORST, PATERNOSTER ROW.

M.DCCC.XLVII.

PLATE XXXV.

FIG.

- 1, 2. *TURBINOLIA MILLETIANA*, p. 196, of the natural size.
3. *TURBINOLIA MILLETIANA*, no. 2, magnified.
4. The coral of *CARYOPHYLLIA SMITHII*, p. 198, of the natural size. From a specimen dredged in Oban Bay, and given to me by Mr. Jos. Alder.
5. A magnified view of the animal of *Caryophyllia Smithii*.
6. The mouth more highly magnified. This, and the preceding figure are from drawings, of my friend Mr. Alder.
7. *CARYOPHYLLIA SMITHII*, dredged in 80 fathoms, off Foulness. From a drawing of Professor Edw. Forbes, who also furnished the following: viz.
8. A tentaculum of *Caryophyllia Smithii*.
9. *ZOANTHUS COUCHII*, p. 202. Copied from Couch's Cornish Fauna, pl. 15, fig. 3.
10. *CORYNACTIS VIRIDIS*, p. 205.
11. A magnified view of the oral disk. These figures of *Corynactis* are engraved from the drawings of Mr. Peach.

\* \* \* Mr. MacAndrew and Professor Forbes have recently again dredged up the *Turbinolia milletiana* off the Cornish coast. "It lives stuck in sand at a depth from 20 to 25 fathoms. We have taken many dead specimens, but only one alive. That, however, is in fine condition. The tentacula are in two rows, of about 24 in each circle, and have globular tips. The mouth is placed across the narrow diameter. Colours very pale." *E. Forbes.*

PLATE XXIV.

